

The relationship between ventricular size and visual function in children with hydrocephalus

T.O.Odebode¹, M.T. Shokunbi^{1,2}, A.O.Malomo^{1,2}, J.B.Familusi³, A.M. Agbeja-Baiyerolu⁴ and A.O. Ogunseyinde⁵
Department of Surgery¹, Anatomy², Paediatrics³, Ophthalmology⁴ and Radiology⁵, University College Hospital, and College of Medicine, University of Ibadan, Ibadan, Nigeria.

Summary

We examined the relationship between ventricular size and visual function in 50 children (36 males and 14 females) with hydrocephalus. The third and lateral ventricular sizes and the visual function scores did not significantly differ between the sexes. Ventricular enlargement was most pronounced at the trigones and least at the level of the foraminae of Monro. The visual function score correlated inversely with the size of the lateral ventricle measured at the levels of the anterior horn and the trigone and expressed as coronal diameter and "Modified" Evans' ratio ($r = 0.49$; $P = 0.001$ and $r = -0.38$, $P = 0.01$ for the anterior horn; $r = 0.48$, $P = 0.001$ and $r = -0.35$, $P = 0.001$ for the trigone). The size of the third ventricle did not correlate with visual function score. A "modified" anterior Evans ratio of 0.60 and a trigonal Evans ratio of 0.73 were associated with very low visual function score. Furthermore, there was significant inverse correlation between occipitofrontal circumference (OFC) and visual function ($r = -0.6379$, $P = 0.001$), but OFC was not valuable for predicting visual function before the onset of head enlargement.

Keywords: *Hydrocephalus; vision; ventricular size; visual function score; children.*

Résumé

Nous avons examiné la relation entre la taille ventriculaire et la fonction visuelle chez 50 enfants (36 mâles et 14 femelles) ayant un hydrocéphale. Les tailles des troisièmes et des ventricules latéraux et la fonction visuelle n'ont pas différencié de manière significative dans les 2 sexes.

L'élargissement ventriculaire n'a pas été prononcé aux tréfontes visuelles. L'élargissement a été inversement proportionnel à la taille des ventricules latéraux, au niveau de la corne antérieure et du trigone. Les valeurs ont été exprimées en fonction du diamètre coronal et du ratio Modifié d'Evans ($r = 0.49$; $P = 0.001$ et $r = 0.38$; $P = 0.01$) pour la corne antérieure; $r = -0.48$, $p = 0.001$ et $r = -0.35$, $p = 0.001$ pour le trigone). La taille du troisième ventricule n'a pas corrélation avec la fonction visuelle. Une modification du ratio antérieur d'Evans (ratio de 0,60) et du trigonal d'Evans (ratio de 0,73) ont été associés à une faible fonction visuelle.

De plus, il y a eu une corrélation inversement significative entre la circonférence occipito-frontale (OFC) et la fonction visuelle ($r = -0,6379$, $P = 0,001$) mais l'OFC n'a pas été valable pour la prédiction de la fonction visuelle, avant le début de l'élargissement de la tête.

Introduction

Hydrocephalus, the commonest congenital malformation of the central nervous system (CNS) seen in Ibadan constitutes 40.6% of all CNS anomalies [1]. Fifty-three percent of affected children present in advanced stages with gross head enlargement and neurological disability including seizures and loss of vision [2]. Visual loss is commonly attributed to optic atrophy resulting from a variety of causes such as papilloedema, chiasmal traction, papillomacular bundle atrophy, developmental abnormalities and third ventricular dilatation acting either singly or in combination [3,4].

One of the consequences of hydrocephalus is that it impairs tissue metabolism by tampering both globally and focally with the cerebral microcirculation [5]. Cortical maturation and myelination, two developmental prerequisites for normal neurological function, are highly dependent on a normal circulation. It is therefore not surprising that a variety of neurobehavioural deficits accompany hydrocephalus. An inverse relationship between ventricular size and intellectual development has been described in infants with myelomeningocele [6]. Furthermore, the neurologic outcome following shunting procedures in advanced hydrocephalus is poor [2].

The precise relationship between ventricular dilatation and visual deterioration in hydrocephalic children is not known. The purpose of this study is to determine first, whether there are any relationship between the extent to which visual function is impaired, and the degree of ventricular dilatation; and, second, to predict the ventricular size below which shunting procedures will produce the best outcome for vision.

Patients and methods

This was a prospective study of fifty consecutive hydrocephalic young children aged 18 months or less, evaluated at our Neurosurgical Clinic between January 1995 and June 1996. Only those who met the study criteria were selected. Infants with closed fontanelles, associated encephalocele or visual impairment traceable to an existing eye disease, were excluded.

The initial evaluation consisted of history and examination to obtain a clinical diagnosis. This was followed by transfontanelle ultrasound scanning using the ACCUOUSON scanner (ATL, Advanced Technology Laboratories Ltd. USA) a B-Mode real-time scan with gray scale imaging. Each patient was scanned in supine position with or without sedation, employing all possible

views with a 7.5 MHz sector transducer, moved to different positions on the scalp or tilted at different angles to obtain corresponding views [6]. An aqueous gel was used in all instances for good skull contact and the scanning was done through the anterior fontanelle. Three examiners scanned all patients. The value for each patient was taken as the mean of the three values obtained. Coronal widths of the third and lateral ventricles were measured. Lateral ventricular width was measured at three levels, i.e., anterior horn, Foramen or Monro and the trigones. Lateral ventricular size was also assessed at the levels of anterior horns and trigones employing the Evans' ratio method [7,8] (described in fig. 1). modified for ultrasonography. The original description of the method was based on computerized tomography [6,8].

Visual function was scored by assessing the following parameters;

- a) Visual acuity, expressed by visual tracking and fixation, and optokinetic eye movements.
- b) Pupillary size and reaction to direct light; and
- c) The severity and extent of optic disc pallor.

Each eye was examined in turn, the other eye having been covered with an occlusive patch. The optic disc was examined after pupillary dilation with 0.5% tropicamide eye drops. Each parameter was scored as shown in table 2. The visual function score for each eye ranged between a minimum of 5 and a maximum of 15 units. The patient's total visual function score (TVFS) was the sum of the scores for both eyes. The range of scores was 10 – 30. The relationship between ventricular size and visual function score was assessed by correlation analysis and standard test of significance. A P value less than 0.05 was taken as significant.

Results

Patients' characteristics

Of the 50 infants studied, 36 were males and 14 were females. At presentation, they ranged in age from one day to eighteen months with a mean of 4.5 ± 2.6 months. Table 1 summarizes their age and sex distribution. More than half of these infants (54%) presented before or at three months of age. Progressive head enlargements, the commonest mode of presentation, occurred in 39 (78%) patients. The remaining infants presented primarily with lumbar myelomeningocele. In this latter group, hydrocephalus was manifest at presentation or was detected by ultrasonography.

Table 1: Age and sex distribution of hydrocephalic children

Age range (mo)*	Males (%)	Females (%)	Total (%)
0 – 3.0	5 (10)	7 (14)	27 (54)
3.1 – 6.0	5 (10)	4 (8)	9 (18)
6.1 – 9.0	7 (14)	3 (6)	10 (20)
9.1 – 12.0	2 (4)	0 (0)	2 (4)
12.1 – 15.0	1 (2)	0 (0)	1 (2)
15.1 – 18.0	1 (2)	0 (0)	1 (2)
Total	36 (72)	14 (28)	50 (100)

Mnths = months




Fig. 1: The lateral and third ventricles as visualised by a coronal Transfontanellear ultrasound scan; and a line diagram of the horizontal section of the brain (showing the lateral and third ventricles) as used for the estimation of Evan's ratio. A = greatest transverse diameter of the cerebrum; B = greatest distance between the anterior horns; C = greatest distance between the trigones; RLV = right lateral ventricle; LLV = left lateral ventricle. Anterior Evan's ratio = B/A; trigonal Evan's ratio = C/A.

Total visual function score (TVFS)

Each of the patient's eyes was scored for visual acuity, pupillary size and reaction to direct light as well as optic disc paor (Table 2). The sum of the scores for both eyes is taken as the total visual function score (TVFS) for the child. The mean \pm SD TVFS for the entire sample was 23.6 ± 5.7 . There was no significant difference between the mean TVFS of males, 23.6 ± 6.0 and females 24.6 ± 4.7 , ($P > 0.05$).

Table 2: Visual function scale (TVFS)

Visual function	Score
Visual fixation and tracking	
Immediate	3
Slow	2
Absent	1
Pupillary size and reaction	
Size less than 4 mm	2
Size 4 mm and above	1
Brisk reaction	3
Sluggish reaction	2
No reaction	1
Optic disk pallor	
Nil	3
Moderate	2
Severe	1
< 25% of disc surface	4
25 – 50% of disc surface	3
50 – 75% of disc surface	2
> 75% of disc surface	1

The visual function score for each eye is the sum of the scores for all parameters. The total visual function score (TVFS) for the patient is the sum of the scores for both eyes
 Minimum score is 10
 Maximum score is 30

Ventricular size (transverse diameters) and visual function

Mean lateral ventricular widths (i.e. transverse diameters) were measured at the level of foramen of Monro, the anterior horns and the trigones and their relationship to visual function was investigated. A significant inverse correlation was found between TVFS and the ventricular diameter at the anterior horns as well as the trigones ($r = 0.49$ at $P = 0.0003$ and $r = -0.48$ at $P = 0.0004$, figures 2a and 2d, respectively).




Fig. 2a: Scatter diagram to illustrate the correlation between TVFS and the transverse diameter of the lateral ventricle at the level of the anterior horn.




Fig. 2b: Scatter diagram to illustrate the correlation between TVFS and the size of the third ventricle.

There were a poor correlation between TVFS on the one hand and the third ventricular width and the size of the lateral ventricle at the level of foramen of Monro (LVZ) on the other ($r = 0.2084$ at $P = 0.1463$ and $r = -0.1411$ at $P = 0.3283$; Fig. 2b and 2c respectively).




Fig. 2c: Scatter diagram to illustrate the correlation between TVFS and the transverse diameter of the lateral ventricle at the level of the foramen of Monro.




Fig. 2d: Scatter diagram to illustrate the correlation between TVFS and the transverse diameter of the lateral ventricle at the level of the trigone.

Ventricular size (modified Evans' ratio) and visual function

a. Modified Anterior Evans' Ratio (mAER) and Visual Function

The mean value of the mAER obtained for the sample was 0.5968 ± 0.1775 . The mean value for males and females were 0.6086 ± 0.1858 and 0.5664 ± 0.1561 , respectively. There was no significant sex difference ($P > 0.05$). The total visual function score (TVFS) declined steadily with increasing mAER. These two parameters, TVFS and mAER, were significantly inversely correlated ($r = -0.3893$ and $P = 0.0052$, Fig. 3a)

b. Modified Trigonal Evans' Ratio (mTER) and Visual Function

The mean lateral ventricular size measured by Evans' ratio at the level of the trigones was 0.7402 ± 0.1697 for the entire sample, 0.7097 ± 0.1544 for males and 0.6900 ± 0.2101 for females. There was no significant sex difference ($P > 0.05$). However, there was a significant inverse correlation total visual function score between and mTER ($r = -0.34631$ and $P = 0.0138$, Fig. 3b)




Fig. 3a: Scatter diagram to illustrate the correlation between TVFS and the modified anterior Evan's ratio.




Fig. 3b: Scatter diagram to illustrate the correlation between TVFS and modified trigonal Evan's ratio.

Ventricular size and severity of visual impairment

Impairment of visual function was empirically categorized into severe, moderate and mild forms based on visual score ranges of 10 – 17, 18 – 26 and 27 – 30 units, respectively. Mean values of mAER and mTER for patients in each category was estimated.

Table 3 shows that 48% of the infants had visual impairment of moderate degree. Infants manifesting severe visual impairment comprised 16% of the group while a mild loss occurred in 36% of the sample. Table 3 also reveals that mean Evans' ratio decrease with decreasing severity of visual impairment and that a mean mAER greater than 0.60 and mTER higher than 0.73 appear to be predictive of imminent severe visual deterioration.




Fig. 3c: Scatter diagram to illustrate the correlation between TVFS and the occipito-frontal circumference.

Occipito-frontal circumference (OFC) and visual function

The maximum measurable occipito frontal circumference (OFC) was obtained using a tape measure in all infants at initial evaluation. Values varied between 36 cm and 68 cm, (mean = 47.06 ± 7.69 cm). OFC greater than 97th percentile for age and sex (macrocephaly) was obtained in 39 (78%) of the 50 infants. In the remaining patients (22%), ventriculomegaly was detected by transfontanelle ultrasound performed because of pre-existing spinal dysraphism. A high inverse correlation was drawn between (OFC) and total visual function score among these infants (Fig. 3c). Comparing other parameters related to visual function in this work, OFC correlated best with visual function (r = -0.65 at P = 0.001).

Discussion

In cases of hydrocephalus, ventricular dilation does not merely represent a progressive accumulation of cerebrospinal fluid due to imbalance between production and absorption, but the absorptive defect sets off a complex interplay between altered intracranial pressure and the biochemical properties of the brain itself. Whether the ventricular dilatation exerts direct influence on cerebral functions has been the subject of numerous previous investigations. The effect of ventricular dilatation on visual function is the subject of the present study.

The study sample consisted of 36 males and 14 females. The male predominance (ratio 2.57:1) confirmed earlier reports of higher incidence of congenital hydrocephalus among males [12,18], although in Binitie's series from northern Nigeria, the sexes were equally affected [13]. Males are known to be at a greater risk than females of developing congenital hydrocephalus [9] and in 2 out of 3 patients with congenital hydrocephalus a lesion obstructing the aqueduct can be demonstrated [14], suggesting that aqueductal stenosis is common among males.

Our results corroborate this finding. Sixty-five percent of the 20 children who had definitive ultrasound diagnosis demonstrated features of aqueductal obstruction, and 85% of these were males. It has also been suggested that in hydrocephalus, males are prone to greater ventricular dilatation than females [15]. Our results do not support this assertion. It is not surprising that ventricular dilatation was most pronounced at the trigones.

Anatomically, the collateral trigones constitute the largest and probably the most distensible portion of the lateral ventricular system [16].

A semiquantitative system of assessment of visual function in pre-literate hydrocephalic infants was pre-literate designed for this study. This system grades the three visual parameters: visual acuity, optic disc palor and pupillary size and reaction to direct light and scores each eye 5-15 units. It's repeatability, reproducibility, rate of agreement by two observers as well as limitations have been subjected to statistical tests (unpublished data).

When total visual function scores obtained using this system were correlated with ventricular size measured at two levels (anterior horns and trigones) using two techniques (coronal diameters and modified Evans' ratio), significant inverse correlations were obtained as follows, in descending order of correlation

(r): anterior coronal width $r = -0.4900$; trigonal coronal width $r = -0.4801$; modified anterior Evans' ratio (mAER) $r = -0.3889$; and modified trigonal Evans' ratio (mTER) $r = -0.3463$.

The third ventricular size did not correlate with visual function score, although correlations between visual function score and Evans' ratio are lower than similar values for coronal diameters, the former is still considered a better indirect predictor of cerebral function than the latter because it incorporates both ventricular and brain parenchyma sizes and reflects not only the extent of ventricular enlargement but also that of cerebral compression when present.

The correlation obtained between ventricular sizes and visual function scores cannot be explained entirely on the basis of anatomic proximity. There was no correlation between the size of the third ventricle (a close relation of the optic chiasma) and visual function. There was a good inverse correlation of visual function score with the anterior horns which maintain no direct contact with the visual pathways. It is easier to explain trigonal correlation based on proximity of the trigones to the optic radiation and close relation between its extension into adjacent horns and the lateral geniculate bodies as well as the calcarine cortex [16].

Clearly, other factors are involved. Cortical histology, cerebral fluid and electrolyte content, intraventricular pressure, cerebral blood flow, oxygen and glucose metabolism over extensive areas of the brain including the visual cortices, and cerebral vascular architecture are altered in hydrocephalus. Their relationship to visual function await further investigation [5, 17, 18, 19].

We graded group means of Evans' ratio and compared them to visual loss graded as severe, moderate and mild. We found mAER greater than 0.60 mTER greater than 0.73 to be indicative of severe visual impairment. It is suggested that infants with such or greater ratios be shunted promptly even when the head size is normal. This is of particular importance in preterm infants and infants with myelomeningocele undergoing serial sonographic screening for follow up of hydrocephalus. It is expected that such prompt intervention will produce a good outcome for visual function.

Excessive enlargement of the head, especially an excessive rate of cranial enlargement, is usually the first and most reliable feature of hydrocephalus in infancy [20]. An occipito-frontal circumference (OFC) greater than 97 percentile for age and sex is highly suggestive of hydrocephalus [21]. This range of OFC was found in 78% of our patients, the largest being 68 cm and smallest 33 cm. Visual scoring revealed a high inverse correlation $r = 0.6379$, $P = 0.001$ between OFC and visual function. Compared with other measured parameters employed in this work, OFC correlated best with visual function.

However, since increase in OFC (i.e., onset of head enlargement) is a late event for hydrocephalus, OFC is considered to be of limited value for predicting early visual deterioration. Although Milhorat (1979) expressed the view that no definite correlation exists between the degree of ventriculomegaly, the size of the head and the eventual development of mental and motor skills [20], our study suggests that OFC is a good but late predictor of visual function after the onset of head

enlargement. OFC is the simplest measurement that predicts visual status even at initial clinical assessment.

References

1. Adelaye A and Odeku E.L. Congenital malformations of the central nervous system in Nigeria. *West Afri Med J* 1972; 21: 73-77.
2. Afolabi A.O and Shokunbi M.T. Socio-economic implications of the surgical treatment of hydrocephalus. *Nigerian J Paed.* 1993; 20: 94-97.
3. Smith J.H, Walsh T.J and Shipley T. Blindness in congenital hydrocephalus. *Am J of Ophthalmology* 1996; 62: 251-257.
4. Duke-Elders S. (ed). *System of ophthalmology. Normal and abnormal development Vol. 3 Part 2 (congenital deformities)*. London, Henry Kimpton 1964; 1078 - 1080.
5. Donahue S. A relationship between fine structure and function of blood vessels in the central nervous system of rabbit foetuses. *Am J Anat* 1964; 115: 17-26.
6. Masato N and Yuh T. Effect of ventricular size of intellectual development in children with myelomeningocele. In: Matsumoto S and Tamaki N (eds) *Hydrocephalus; pathogenesis and treatment*. Tokyo Springer - Verlag 1991; 563 - 570.
7. Mimi Magio. In: Saunders RC (ed.) *Clinical sonography, a practical guide, 2nd ed.* Boston Little Brown and Co. 1991: 421 - 436.
8. Yamada H. *Pediatric cranial computed tomography - Igaku - Shoin, Tokyo, 1983: 155 - 120.*
9. Chapman P.H. Hydrocephalus in childhood. In Youmans JR (ed.) *Neurosurgical surgery Vol 2. 3rd ed.* Philadelphia WB Saunders Co 1990: 1236 - 1276.
10. Storrs, B.B. Ventricular size and intelligence in myelodysplastic children. In: Marlin AE (ed.) *Concepts in paediatric neurosurgery, Vol. 8.* Karger, Basel 1998: 51-56.
11. Raimondi A.J and Soare P. Intellectual development in shunted hydrocephalic children *Am J Dis Child* 1974; 27: 664 - 671.
12. Gupta B. **Incidence of congenital malformations in Nigerian children.** *West Afr Med J* 1969; 18: 22 - 27.
13. Binitie OP. Congenital malformations of the central nervous system at Jos University Teaching Hospital, Jos (Nigeria) *West Afri Med J* 1992; 11: 7-12.
14. Mori K. Hydrocephalus. Revision of its Classification In: Matsumoto S and Tamaki N (eds.) *Hydrocephalus; pathogenesis and treatment*, Tokyo Springer - Verlag. 1991; 361 - 368.
15. Margaret D.J. Growth and development survey (Unpublished communication) Institute of Child Health. University of Ibadan, Ibadan, Nigeria. 1980: 1-2.
16. Carpenter B.M. *Core Text of Neuroanatomy, 4th ed.* Baltimore. Williams and Wilkins Co. 1991: 9 - 27, 39 - 42 and 46 - 47.
17. Sato H, Sato N, Tamaki N, Matsumoto S. Cerebral haemodynamics in hydrocephalus during infancy: experimental and clinical

- studies. In: Matsumoto S and Tamaki N (eds) *Hydrocephalus: Pathogenesis and treatment* Tokyo: Springer Verlag 1991: 131 – 143.
18. Greitz YUB, Grope AOL, Kalmer MSF and Lopez J. Pre and post operative evaluation of cerebral blood flow in low pressure hydrocephalus *N Neurosurg* 1969; 31: 644 – 651.
19. Shirane R, Sato S, Kameyama M *et al.* Cerebral blood flow and oxygen metabolism in children with hydrocephalus. In: Matsumoto S and Tamaki N. (eds). *Hydrocephalus; Pathogenesis and treatment*. Tokyo Springer – Verlag. 1991: 144 – 152.
20. Milhorat TH. Treatment of hydrocephalus in infancy. In: Lindsay Symon. (ed) *Operative surgery. Fundamental international techniques. Neurosurgery*, London. Buterworths, 197: 388 – 394.
21. Clain A. (ed). *Hamilton Bailey's demonstration of physical signs in clinical surgery*. 17th ed. Bristol ELBS. John Wright and Sons Ltd. 1986: 54 – 56.